



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

264-1204

Date of Issuance:

10/9/18

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Luxxur A Herbicide

Bayer CropScience  
 2 T.W. Alexander Dr.  
 Research Triangle Park, NC 27709

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 264-1204."
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Signature of Approving Official:

Date:

10/9/18

Shaja Joyner, Product Manager 20  
 Fungicide-Herbicide Branch, Registration Division (7505P)

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 10/02/2018

If you have any questions, please contact Kathryn Montague by phone at 703-305-1243, or via email at [montague.kathryn@epa.gov](mailto:montague.kathryn@epa.gov)

Enclosure

**ACCEPTED**

10/09/2018

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 264-1204

TRIBENURON METHYL GROUP 2 HERBICIDE

# LUXXUR A HERBICIDE

For: Use on Wheat (Spring, Winter, Durum)

**ACTIVE INGREDIENT:**

Tribenuron methyl

Methyl 2-[[[4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate..... 50.00%

**OTHER INGREDIENTS:** ..... 50.00%

**TOTAL:** ..... 100.00%

EPA Reg. No. 264-XXXX

EPA Est.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Please refer to [back panel] [booklet] for additional precautionary statements and directions for use. [Note to reviewer: Location of additional precautionary statements and directions for use will vary between those listed, depending on container type/size.]

Net Contents:

**PRODUCED FOR**



Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

## FIRST AID

### If on Skin:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

- Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.
- Avoid contact with skin, eyes or clothing.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.
- Shoes plus socks

### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

---

## CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

---

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

---

## DIRECTIONS FOR USE

---

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
Read the entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval **(REI) of 12 hours**.

**PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:**

- Coveralls
- Chemical resistant gloves made of any waterproof material.
- Shoes plus socks.

## PRODUCT INFORMATION

### LUXXUR A HERBICIDE:

- provides control of certain broad-leaved weeds in wheat (spring, winter and durum). Warm, moist, growing conditions promote active weed growth and enhance the activity of LUXXUR A HERBICIDE by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress such as cold weather, drought stress or excessive heat may not be adequately controlled or suppressed, and regrowth may occur. For best results, ensure thorough spray coverage of target weeds.
- Rapidly stops growth of susceptible weeds through absorption of the foliage. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application.

## USE RESTRICTIONS

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not discharge excess material on the soil at a single spot in the field, or mixing/loading station.
- Do not apply LUXXUR A HERBICIDE to wheat that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- Do not store pesticides near well sites.
- Do not apply to wheat underseeded with another crop.
- Pre-Harvest Interval (PHI): Do not harvest within 45 days of the last application.
- Do not apply LUXXUR A HERBICIDE within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment because crop injury may result.
- Do not use LUXXUR A HERBICIDE plus Malathion because crop injury may result.
- Do not apply LUXXUR A HERBICIDE by air in the state of New York.
- LUXXUR A HERBICIDE is not for use in the state of California.

### Grazing, Feeding, and Harvesting

- Allow at least 7 days between application and grazing of treated forage.
- Allow at least 7 days between application and feeding of forage (green chop) from treated areas to livestock.
- Allow at least 30 days between application and feeding of hay from treated areas to livestock.
- Allow at least 45 days between application and harvesting of grain. Harvested straw may be used for bedding and/or feed.

When using LUXXUR A HERBICIDE in tank mixes or sequential applications with other products containing tribenuronmethyl, do not exceed the following limits:

Active Ingredient: Tribenuron-methyl						
Use	Maximum AI Oz/A per Single Application	Maximum Oz/A of Product per Single Application	Maximum AI Oz/A per Use Period*	Maximum Oz/A of Product per Use Period*	Maximum Number of Applications per Use Period*	Pre-Harvest Interval, Days
Wheat	0.25	0.50	0.25	0.50	2	45 (for grain)

\* Use Period:  
One maximum application per year or two applications not to exceed the maximum rate per year, see above chart for specific crop restrictions.

## USE PRECAUTIONS

- Injury or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:
- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than those listed on this label.
- Varieties of wheat (including durum) may differ in their response to various herbicides. Consult your state experiment station, university, or extension agent as to crop sensitivity of any herbicide. If no information is available, limit the initial use to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after LUXXUR A HERBICIDE application, temporary discoloration and/or crop injury may occur.
- Dry, dusty field conditions may result in reduced control in wheel track areas.
- Calibrate sprayers only with clean water away from well sites.
- Make scheduled checks of spray equipment.

- Mix only enough product for the job at hand.
- Ensure that all operation employees accurately measure pesticides.
- Avoid overfilling of spray tank.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

## APPLICATION INSTRUCTIONS

### Aerial Application

- For aerial application, select nozzles and pressure that deliver medium or coarse spray (ASABE Standard S572.1) and that provide optimum spray distribution and maximum coverage at a minimum of 5 gal/A (GPA).
- Do not apply LUXXUR A HERBICIDE by air in the state of New York.
- For aerial applications, do not apply during a temperature inversion, when wind speed is less than 3 mph or above 10 mph, or when conditions favor poor coverage and/or off-target spray drift.

### Ground Application

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

- Select nozzles and pressure that deliver medium spray droplets.
- Nozzles that deliver coarse spray droplets may be used to reduce drift, provided spray volume is increased to maintain coverage on small weeds. For optimal product performance and minimal spray drift, adjust the spray boom to the lowest possible spray height listed in manufacturers' specifications.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.
- For flat-fan nozzles, use a spray volume of at least 10 GPA.
- For flood nozzles on 30" spacing, use flood nozzles no larger than TK 10 (or the equivalent), a pressure of at least 30 psi and a spray volume of at least 10 GPA. For 40" nozzle spacing, use at least 13 GPA. For 60" spacing, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.
- Use screens that are 50-mesh or larger (metal or nylon).

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather related factors determine the potential of spray drift. The applicator is responsible for considering all of these factors when making application decisions. Consult the local Cooperative Extension for additional information. Avoiding spray drift is the responsibility of the applicator.

### DROPLET SIZE

To obtain uniform spray coverage and reduce spray drift, equip sprayers with nozzles that provide medium to coarse spray droplets as defined in ASABE Standard S572.1.

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions! When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry. Avoid spraying during conditions of low humidity and/or high temperatures. See **Wind, Temperature Inversions, and Sensitive Areas** sections of this label.

### WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. Applications during gusty or calm wind conditions should be avoided. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. For applications made in-furrow or below soil-level, wind speed restrictions are not applicable.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### TEMPERATURE INVERSIONS

Drift potential is high during temperature inversions and applications should be avoided under these conditions. Temperature inversions are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. If fog is not present, inversions can also be identified by the movement of smoke or dust from a ground source -- smoke or dust that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion.

## **SENSITIVE AREAS**

When applying adjacent to residential areas, water bodies, habitats known to have threatened or endangered species, or non-target crops, drift can be minimized to these areas by making application when the wind direction is away from these areas.

### **Aerial Applications**

- Mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices.
- The minimum practical boom length should be used, and should not exceed 75% of the wing span or rotor diameter.
- Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety.

### **Ground Application**

- Wind Speed: Avoid spraying when sustained wind speeds approach or exceed 10 mph. Avoid applications in gusty wind conditions.

## **HERBICIDE RESISTANCE MANAGEMENT (WSSA) RECOMMENDATIONS**

For resistance management, LUXXUR A HERBICIDE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to LUXXUR A HERBICIDE and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of LUXXUR A HERBICIDE or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage ( or other mechanical control methods), cultural ( e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

# SPECIFIC USE DIRECTIONS

## FOR WEED CONTROL IN WHEAT (SPRING, DURUM AND WINTER)

### LUXXUR A HERBICIDE + LUXXUR B HERBICIDE

Tank Mix Partner	Product Rates fl oz/A	WEEDS CONTROLLED	GROWTH STAGE TIMING
<b>LUXXUR A HERBICIDE + LUXXUR B HERBICIDE</b>	<b>0.214 + 6.85</b>	<b>Weeds controlled by LUXXUR B HERBICIDE alone +</b> Canada thistle*, Perennial sowthistle*, Dandelion, Scentless chamomile, Narrow-leaved hawk's-beard, Lamb's quarter, Cow cockle	Wheat (spring and durum) may be treated from 2 to 6 leaf stage on main stem, plus 3 tillers, but prior to jointing (presence of first node).  Winter wheat may be treated when the majority of plants have two leaves to full tillering, but prior to jointing (presence of first node).  Weed growth stages: <ul style="list-style-type: none"> <li>• Canada thistle and Perennial Sowthistle: up to 6 inches in height.</li> <li>• Dandelion: up to 8 inches in diameter;</li> <li>• Scentless chamomile: up to 4 inches in height;</li> <li>• Narrow-leaved hawk's-beard: prior to bolting;</li> <li>• Lamb's quarter and cow cockle: up to 4 inches in height.</li> </ul>
*Season long control, with some regrowth in the fall.			

### Tank Mix Instructions

For control of weeds not listed on this label, Luxxur A Herbicide may be mixed with other herbicides with the exception of dicamba containing products. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Broadleaf Herbicides			
MCPA Ester <sup>1, 3, 4</sup>	Starane <sup>®2,3</sup> / Starane Ultra <sup>2,3</sup>	2,4-D Ester <sup>1,4</sup>	Sentrallas <sup>®</sup>
<sup>1</sup> MCPA Ester and 2,4-D Ester may be added alone, but not as a component of a premix. <sup>2</sup> Equivalent fluroxypyr product may be substituted. <sup>3</sup> Applications of Luxxur A Herbicide in winter, spring or durum wheat in combination with these EC tankmix partners or equivalents may cause crop response. <sup>4</sup> MCPA Ester and 2,4-D Ester may be added at a rate of no more than 0.25 lb ai/A.			

Grass Herbicides			
Olympus <sup>1</sup>	Osprey	Puma <sup>2</sup>	Rimfire Max
<sup>1</sup> Olympus can be added. Refer to Olympus label concerning crop rotation restrictions. <sup>2</sup> Equivalent fenoxaprop-based products may be substituted			

### Tank Mixtures for Disease Control

- This product may be tank mixed or used sequentially with fungicides registered for use on wheat.

### Tank Mixtures for Insect Control

- LUXXUR A HERBICIDE may be tank mixed or used sequentially with insecticides registered for use on wheat. However, under certain conditions (drought stress, or if the crop is in the 2-4 leaf stage), tank mixes of LUXXUR A HERBICIDE with organophosphate insecticides (such as Lorsban) may produce temporary crop yellowing, or, in severe cases, crop injury. The

potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

#### USE RESTRICTIONS

- Do not apply LUXXUR A HERBICIDE within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment as crop injury may result.
- Do not use LUXXUR A HERBICIDE with malathion because crop injury may result.

#### MIXING INSTRUCTIONS

1. Add one-third of the necessary volume of water to the spray tank.
2. With the agitator running, add the required amount of LUXXUR A HERBICIDE. Ensure that the herbicide is completely dissolved before adding other components of the tank mix.
3. When LUXXUR A HERBICIDE is completely dissolved, add the required amount of the appropriate tank mix partners.
4. Specific Tank Mix Instructions:  
LUXXUR A HERBICIDE + LUXXUR B HERBICIDE: once LUXXUR A HERBICIDE is completely dissolved, add the required amount of LUXXUR B HERBICIDE, then add required amount of non-ionic surfactant or ammonium sulphate last.
5. Add the rest of the water. If an antifoam agent is required, add this last.
6. Use a minimum spray volume of 5 GPA for aerial and 10 GPA for ground applications.
7. Fill tank with fresh water as described in step 1, and continue as directed. Emulsifiable concentrates may make it more difficult to dissolve LUXXUR A HERBICIDE
8. If the spray preparation is left standing without agitation, thoroughly agitate to resuspend the tank mixture before spraying. Use spray preparation of LUXXUR A HERBICIDE Tank Mix within 24 hours or product degradation may occur.

#### MINIMUM RECROPPING INTERVAL AND CROP ROTATION GUIDELINES

Observe the rotational cropping restrictions stated on the LUXXUR B HERBICIDE label.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide storage

Store in a cool, dry place.

#### Pesticide disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Container handling

**Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.**

#### **Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container.**

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

#### **Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container.**

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

#### **Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container.**

Do not reuse or refill this container.

Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure

that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container.**

Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**Refillable Fiber Drums With Liners: Refillable container (fiber drum only).**

Refilling Fiber Drum: Refill this fiber drum with LUXXUR A HERBICIDE containing tribenuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**All Other Refillable Containers: Refillable container.**

Refilling Container: Refill this container with LUXXUR A HERBICIDE containing tribenuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact FMC at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container.**

Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

**LUXXUR A HERBICIDE (PENDING) 10/2/2018, 10/5/2018**